


EMD Protocol 13

ALL ANSWERS CAN BE FOUND ON THE PROTOCOL



www.CrosswordWeaver.com

ACROSS

- 3 A state of unconsciousness from which the patient cannot be aroused
- 5 EMDs should not advise administration of oral sugar to ___ diabetics
- 9 Hypoglycemia is commonly confused with alcohol ___
- 10 Stay on the line with caller if her/his condition seems ___ or is worsening
- 16 Insulin Shock
- 19 Diabetes is a ___ that EMDs may accept at face value because of its high degree of accuracy
- 20 Determining the level of ___ is the key to correctly assigning the prehospital response
- 21 13-D-1

DOWN

- 1 Is s/he ___ alert
- 2 Airway control is the first ___ in PDIs if the patient is unconscious
- 4 The ___ of an unconscious patient must be constantly maintained
- 6 Diabetic Coma (later ___)
- 7 Diabetic ___ (gradual onset)
- 8 An early sign of ___ blood sugar is abnormal behavior
- 11 Suffix "C" represents ___ or aggressive
- 12 An ___ sign of low blood sugar is abnormal behavior
- 13 Is s/he breathing ___
- 14 ___ Shock when too much insulin has depleted the body's available blood sugar
- 15 Is s/he ___ normally now
- 17 The ketoacids (___) produced are toxic to patient and cause a slowly
- 18 Alert ___ behaving normally (13-A-1)